Leader in Biomolecular Solutions for Life Science

# mTOR Rabbit pAb

Catalog No.: A24743 1 Publications



## **Basic Information**

#### **Observed MW**

289kDa

#### **Calculated MW**

289kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB

# **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene belongs to a family of phosphatidylinositol kinase-related kinases. These kinases mediate cellular responses to stresses such as DNA damage and nutrient deprivation. This kinase is a component of two distinct complexes, mTORC1, which controls protein synthesis, cell growth and proliferation, and mTORC2, which is a regulator of the actin cytoskeleton, and promotes cell survival and cell cycle progression. This protein acts as the target for the cell-cycle arrest and immunosuppressive effects of the FKBP12-rapamycin complex. Inhibitors of mTOR are used in organ transplants as immunosuppressants, and are being evaluated for their therapeutic potential in SARS-CoV-2 infections. Mutations in this gene are associated with Smith-Kingsmore syndrome and somatic focal cortical dysplasia type II. The ANGPTL7 gene is located in an intron of this gene.

### **Recommended Dilutions**

WB

1:500 - 1:1000

# **Immunogen Information**

**Gene ID** 

2475

**Swiss Prot** 

P42345

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

#### **Synonyms**

SKS; FRAP; FRAP1; FRAP2; RAFT1; RAPT1; mTOR

# Contact

€

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of lysates from Hep G2 cells using mTOR Rabbit pAb (A24743) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.