

A24909

Leader in Biomolecular Solutions for Life Science



IκBα Rabbit mAb

Catalog No.: A24909

Recombinant

2 Publications

Basic Information

Observed MW

35kDa

Calculated MW

35kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC63567

Background

Enables NF-kappaB binding activity and heat shock protein binding activity. Involved in several processes, including cytoplasmic sequestering of NF-kappaB; leukocyte migration involved in inflammatory response; and liver regeneration. Acts upstream of or within cellular response to cytokine stimulus and cellular response to organic cyclic compound. Located in cytosol and nucleus. Part of protein-containing complex. Used to study gastritis; pulmonary emphysema; and sciatic neuropathy. Biomarker of several diseases, including acute necrotizing pancreatitis; alcoholic hepatitis; artery disease (multiple); middle cerebral artery infarction; and toxic shock syndrome. Human ortholog(s) of this gene implicated in several diseases, including childhood-onset asthma; hematologic cancer (multiple); hepatitis B; hepatitis C; and lung disease (multiple). Orthologous to human NFKBIA (NFKB inhibitor alpha).

Recommended Dilutions

WB 1:2000 - 1:7000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

25493

Swiss Prot

Q63746

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

RL/IF-1; IκBα

Contact

 www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of various lysates, using IkBα Rabbit mAb (A24909) at 1:6000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.