ABclonal®

ABflo® 647 Rabbit anti-Human FAP mAb

Catalog No.: A24966

Basic Information

Observed MW

Calculated MW

27kDa/87kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC64735-ABflo647

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Background

The protein encoded by this gene is a homodimeric integral membrane gelatinase belonging to the serine protease family. It is selectively expressed in reactive stromal fibroblasts of epithelial cancers, granulation tissue of healing wounds, and malignant cells of bone and soft tissue sarcomas. This protein is thought to be involved in the control of fibroblast growth or epithelial-mesenchymal interactions during development, tissue repair, and epithelial carcinogenesis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Immunogen Information

Gene ID2191

Swiss Prot
Q12884

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

FAP; DPPIV; FAPA; FAPalpha; SIMP; prolyl endopeptidase FAP

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.


Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Contact

€

www.abclonal.com

Validation Data





Flow cytometry: 1X10^6 Jurkat cells (negative control,left) and U-87MG cells (right) were surface-stained with ABflo® 647 Rabbit anti-Human FAP mAb (A24966,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 U-87MG cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,left) or ABflo® 647 Rabbit anti-Human FAP mAb (A24966,5 µl/Test,right).