

A24968

Leader in Biomolecular Solutions for Life Science



## ABflo® 488 Rabbit anti-Human CD1d mAb

Catalog No.: A24968

### Basic Information

#### Observed MW

#### Calculated MW

38kDa

#### Category

SMab Recombinant Monoclonal  
Antibody

#### Applications

FC

#### Cross-Reactivity

Human

#### CloneNo number

ARC64590-ABflo488

#### Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

### Recommended Dilutions

FC 5 µl per 10<sup>6</sup> cells in  
100 µl volume

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Background

This gene encodes a divergent member of the CD1 family of transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. The protein encoded by this gene localizes to late endosomes and lysosomes via a tyrosine-based motif in the cytoplasmic tail. Two transcript variants encoding different isoforms have been found for this gene.

### Immunogen Information

#### Gene ID

912

#### Swiss Prot

P15813

#### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### Synonyms

R3; CD1A; R3G1

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification



Affinity purification

#### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Flow cytometry:  $1 \times 10^6$  K-562 cells (negative control, left) and Jurkat cells (right) were surface-stained with ABflo® 488 Rabbit anti-Human CD1d mAb (A24968, 5 µl/Test, orange line) or ABflo® 488 Rabbit IgG isotype control (A22069, 5 µl/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1 \times 10^6$  Jurkat cells were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069, 5 µl/Test, left) or ABflo® 488 Rabbit anti-Human CD1d mAb (A24968, 5 µl/Test, right).