# ABclonal www.abclonal.com

# **SAT1 Rabbit pAb**

Catalog No.: A2506 2 Publications

## **Basic Information**

# **Observed MW**

20-30kDa

#### **Calculated MW**

20kDa

#### **Category**

Polyclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Mouse

# **Background**

The protein encoded by this gene belongs to the acetyltransferase family, and is a ratelimiting enzyme in the catabolic pathway of polyamine metabolism. It catalyzes the acetylation of spermidine and spermine, and is involved in the regulation of the intracellular concentration of polyamines and their transport out of cells. Defects in this gene are associated with keratosis follicularis spinulosa decalvans (KFSD). Alternatively spliced transcripts have been found for this gene.

# **Recommended Dilutions**

WB 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**6303

Swiss Prot
P21673

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

# **Synonyms**

SAT; DC21; KFSD; SSAT; KFSDX; SSAT-1; SAT1

# Contact


www.abclonal.com

### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from Mouse placenta, using SAT1 Rabbit pAb (A2506) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.