# **PSMA4** Rabbit pAb

Catalog No.: A2511



### **Basic Information**

#### **Observed MW**

28kDa

#### **Calculated MW**

29kDa

### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,IP,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a core alpha subunit of the 20S proteosome, which is a highly ordered ring-shaped structure composed of four rings of 28 non-identical subunits. Proteasomes cleave peptides in an ATP- and ubiquitin-dependent manner.

# **Recommended Dilutions**

WB 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

**IP** 0.5μg-4μg antibody for

200µg-400µg extracts

of whole cells

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

### **Contact**

www.abclonal.com

## **Immunogen Information**

 Gene ID
 Swiss Prot

 5685
 P25789

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

# **Synonyms**

HC9; PSC9; HsT17706; PSMA4

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**




Western blot analysis of various lysates using PSMA4 Rabbit pAb (A2511) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of U2OS cells using PSMA4 Rabbit pAb (A2511). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.