

# CD38 Rabbit mAb

Catalog No.: A25398 **Recombinant**

## Basic Information

### Observed MW

45kDa

### Calculated MW

14kDa/34kDa

### Category

SMab Recombinant Monoclonal Antibody

### Applications

WB,IHC-P,FC,ELISA

### Cross-Reactivity

Human

### CloneNo number

ARC66212

## Background

The protein encoded by this gene is a non-lineage-restricted, type II transmembrane glycoprotein that synthesizes and hydrolyzes cyclic adenosine 5'-diphosphate-ribose, an intracellular calcium ion mobilizing messenger. The release of soluble protein and the ability of membrane-bound protein to become internalized indicate both extracellular and intracellular functions for the protein. This protein has an N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites. Crystal structure analysis demonstrates that the functional molecule is a dimer, with the central portion containing the catalytic site. It is used as a prognostic marker for patients with chronic lymphocytic leukemia. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

WB	1:3000 - 1:18000
IHC-P	1:500 - 1:2000
FC	1:500 - 1:1000
ELISA	Recommended starting concentration is 1 $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements.

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Immunogen Information

### Gene ID

952

### Swiss Prot

P28907

### Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### Synonyms

ADPRC1; cADPR1; ADPRC 1

## Product Information



Source	Isotype	Purification
Rabbit	IgG	Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% Sodium azide, 0.05% BSA, 50% glycerol, pH 7.3.


## Validation Data




Immunohistochemistry analysis of paraffin-embedded Human tonsil tissue using CD38 Rabbit mAb (A25398) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human colon tissue using CD38 Rabbit mAb (A25398) at a dilution of 1:800 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.




Flow cytometry: 1X10<sup>6</sup> Hep G2 cells (negative control, left) and Daudi cells (right) were surface-stained with CD38 Rabbit mAb (A25398, 2 µg/mL, orange line) or ABflo® 647 Rabbit IgG isotype control (A22070, 5 µl/Test, blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1X10<sup>6</sup> Daudi cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070, 5 µl/Test, left) or CD38 Rabbit mAb (A25398, 2 µg/mL, right).



Flow cytometry: 1X10<sup>6</sup> Human PBMC were surface-stained with CD38 Rabbit mAb (A25398, 2 µg/mL, orange line) or ABflo® 647 Rabbit IgG isotype control (A22070, 5 µl/Test, blue line), followed by Alexa Fluor® 647 conjugated goat anti-



Flow cytometry: 1X10<sup>6</sup> Human PBMC were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070, 5 µl/Test, left) or CD38 Rabbit mAb (A25398, 2 µg/mL, right).

## Validation Data

---

rabbit pAb staining. Non-fluorescently stained Human PBMC were used as blank control (red line).