

A26342

Leader in Biomolecular Solutions for Life Science



β-Amyloid Rabbit mAb

Catalog No.: A26342

Recombinant

Basic Information

Observed MW

Refer to figures

Calculated MW

87kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

IHC-P, IF/ICC, ELISA, DB

Cross-Reactivity

Mouse

Background

This gene encodes a cell surface receptor and transmembrane precursor protein that is cleaved by secretases to form a number of peptides. Some of these peptides are secreted and can bind to the acetyltransferase complex APBB1/TIP60 to promote transcriptional activation, while others form the protein basis of the amyloid plaques found in the brains of patients with Alzheimer disease. In addition, two of the peptides are antimicrobial peptides, having been shown to have bacteriocidal and antifungal activities. Mutations in this gene have been implicated in autosomal dominant Alzheimer disease and cerebroarterial amyloidosis (cerebral amyloid angiopathy). Multiple transcript variants encoding several different isoforms have been found for this gene.

Recommended Dilutions

DB 1:500 - 1:1000

IHC-P 1:500 - 1:2000

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

www.abclonal.com

Immunogen Information

Gene ID

351

Swiss Prot

P05067

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

AAA; AD1; PN2; ABPP; APPI; CVAP; ABETA; PN-II; preA4; CTFgamma; alpha-sAPP

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data



Dot-blot analysis of N-terminal peptide of β-Amyloid,β-Amyloid 40 proteins,β-Amyloid 42 proteins using β-Amyloid Rabbit mAb (A26342) at 1:1000 dilution.



Immunohistochemistry analysis of paraffin-embedded (5XFAD)Mouse brain and normal mouse brain tissue using β-Amyloid Rabbit mAb (A26342) at a dilution of 1:800 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer (pH 9.0) prior to IHC staining.



Confocal imaging of paraffin-embedded 5XFAD Mouse brain and Mouse brain tissue using β-Amyloid Rabbit mAb (A26342, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Microwave antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IF staining. Objective: 40x.