

A2771

Leader in Biomolecular Solutions for Life Science



Acetyl-Histone H3-K27 Rabbit mAb

Catalog No.: A2771

Recombinant

4 Publications

Basic Information

Observed MW

17kDa

Calculated MW

15kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB,IHC-P,IF/ICC,IP,ChIP,ELISA,DB

Cross-Reactivity

Human,Mouse,Rat,Other (Wide Range
Predicted)

CloneNo number

ARC53670

Recommended Dilutions

WB 1:10000 - 1:120000

DB 1:500 - 1:2000

IHC-P 1:500 - 1:1000

IF/ICC 1:50 - 1:200

IP 0.5µg-4µg antibody for
200µg-400µg extracts
of whole cells

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

ChIP 2µg antibody for
5µg-10µg of Chromatin

Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Contact

 www.abclonal.com


Validation Data




Western blot analysis of lysates from C2C12 cells using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at 1:100000 dilution. C2C12 cells were treated with TSA (1 μ M) at 37°C for 18 hours.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 20 μ g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 30s.




Dot-blot analysis of all sorts of peptides using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at 1:200000 dilution.




Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human thyroid cancer tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human small intestine tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human breast cancer tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human liver tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.


Validation Data




Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse colon tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1:1000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunofluorescence analysis of HeLa treated with TSA and HeLa cells using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunoprecipitation analysis of 600 μg extracts of HeLa cells treated with TSA using 5 μg Acetyl-Histone H3-K27 Rabbit mAb (A2771). Western blot was performed from the immunoprecipitate using Acetyl-Histone H3-K27 antibody (A2771) at a dilution of 1:50000.

Validation Data



Immunoprecipitation of Acetyl-Histone H3-K27 from 600 μ g extracts of NIH/3T3 cells treated with TSA(1 μ M , 18h) was performed using 2 μ g of Acetyl-Histone H3-K27 Rabbit mAb (A2771). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X non-reducing Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Acetyl-Histone H3-K27 Rabbit mAb (A2771) at a dilution of 1 : 50000.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using Acetyl-Histone H3-K27 antibody (A2771) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.