# ABclonal www.abclonal.com

# **NEUROG3 Rabbit pAb**

Catalog No.: A2772 1 Publications

### **Basic Information**

#### **Observed MW**

25kDa

#### **Calculated MW**

23kDa

### Category

Polyclonal Antibody

### **Applications**

WB,IHC-P,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

The protein encoded by this gene is a basic helix-loop-helix (bHLH) transcription factor involved in neurogenesis. The encoded protein likely acts as a heterodimer with another bHLH protein. Defects in this gene are a cause of congenital malabsorptive diarrhea 4 (DIAR4).

# **Recommended Dilutions**

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

**ELISA** Recommended starting

> concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

### **Immunogen Information**

**Gene ID Swiss Prot** 50674 Q9Y4Z2

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

ngn3; Atoh5; NGN-3; Math4B; bHLHa7; NEUROG3

### Contact

0 www.abclonal.com

### **Product Information**


Source **Isotype Purification** Rabbit Affinity purification IgG

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of various lysates, using NEUROG3 Rabbit pAb (A2772) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 


dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Human colon using NEUROG3 Rabbit pAb (A2772) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.