# ABclonal www.abclonal.com

# **DRD1** Rabbit pAb

Catalog No.: A2893 2 Publications

### **Basic Information**

#### **Observed MW**

49kDa

#### **Calculated MW**

49kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Rat

## Background

This gene encodes the D1 subtype of the dopamine receptor. The D1 subtype is the most abundant dopamine receptor in the central nervous system. This G-protein coupled receptor stimulates adenylyl cyclase and activates cyclic AMP-dependent protein kinases. D1 receptors regulate neuronal growth and development, mediate some behavioral responses, and modulate dopamine receptor D2-mediated events. Alternate transcription initiation sites result in two transcript variants of this gene.

## **Recommended Dilutions**

WB 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 1812 P21728

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

DADR; DRD1A; DRD1

### **Contact**

www.abclonal.com €

### **Product Information**


Source **Isotype Purification** Rabbit Affinity purification IgG

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of lysates from Rat brain, using DRD1 Rabbit pAb (A2893) at 1:600 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 0.2s.