# ABclonal www.abclonal.com

## **FST Rabbit pAb**

Catalog No.: A2936

## **Basic Information**

#### **Observed MW**

38kDa

#### **Calculated MW**

38kDa

#### Category

Polyclonal Antibody

## **Applications**

WB,ELISA

## **Cross-Reactivity**

Human, Rat

## **Background**

Follistatin is a single-chain gonadal protein that specifically inhibits follicle-stimulating hormone release. The single FST gene encodes two isoforms, FST317 and FST344 containing 317 and 344 amino acids respectively, resulting from alternative splicing of the precursor mRNA. In a study in which 37 candidate genes were tested for linkage and association with polycystic ovary syndrome (PCOS) or hyperandrogenemia in 150 families, evidence was found for linkage between PCOS and follistatin.

## **Recommended Dilutions**

**WB** 1:100 - 1:500

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

Gene ID Swiss Prot 10468 P19883

## **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

FS: FST

## Contact

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using FST Rabbit pAb (A2936) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.