# ABclonal www.abclonal.com

# **PAR4 Rabbit pAb**

Catalog No.: A3063

#### **Basic Information**

#### **Observed MW**

43kDa

#### **Calculated MW**

37kDa

#### **Category**

Polyclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

### **Background**

This gene encodes a tumor suppressor protein that selectively induces apoptosis in cancer cells through intracellular and extracellular mechanisms. The intracellular mechanism involves the inhibition of pro-survival pathways and the activation of Fasmediated apoptosis, while the extracellular mechanism involves the binding of a secreted form of this protein to glucose regulated protein 78 (GRP78) on the cell surface, which leads to activation of the extrinsic apoptotic pathway. This gene is located on the unstable human chromosomal 12q21 region and is often deleted or mutated different tumors. The encoded protein also plays an important role in the progression of age-related diseases.

#### **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 5074 Q96IZ0

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

#### **Synonyms**

PAR4; Par-4

#### Contact

www.abclonal.com


#### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using PAR4 Rabbit pAb (A3063) at 1:400 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.