

A3736

Leader in Biomolecular Solutions for Life Science



Desmin Rabbit mAb

Catalog No.: A3736

Recombinant

11 Publications

Basic Information

Observed MW

54kDa

Calculated MW

54kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

CloneNo number

ARC0235

Recommended Dilutions

WB 1:1000 - 1:5000

IHC-P 1:1000 - 1:5000

IF/ICC 1:100 - 1:1000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact



www.abclonal.com

Background

This gene encodes a muscle-specific class III intermediate filament. Homopolymers of this protein form a stable intracytoplasmic filamentous network connecting myofibrils to each other and to the plasma membrane. Mutations in this gene are associated with desmin-related myopathy, a familial cardiac and skeletal myopathy (CSM), and with distal myopathies.

Immunogen Information

Gene ID

1674

Swiss Prot

P17661

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

CSM1; CSM2; CDCD3; LGMD1D; LGMD1E; LGMD2R; Desmin

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


Validation Data




Western blot analysis of various lysates using Desmin Rabbit mAb (A3736) at 1:1000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 1s.




Immunohistochemistry analysis of paraffin-embedded Human stomach tissue using Desmin Rabbit mAb (A3736) at a dilution of 1:1600 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human smooth muscle tissue using Desmin Rabbit mAb (A3736) at a dilution of 1:1600 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human prostate cancer tissue using Desmin Rabbit mAb (A3736) at a dilution of 1:1600 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human thyroid tissue using Desmin Rabbit mAb (A3736) at a dilution of 1:1600 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human lung tissue using Desmin Rabbit mAb (A3736) at a dilution of 1:1600 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.



Immunofluorescence analysis of paraffin-embedded mouse heart using Desmin Rabbit mAb (A3736) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Validation Data



Immunofluorescence analysis of paraffin-embedded rat heart using Desmin Rabbit mAb (A3736) at dilution of 1:100 (40x lens).
Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.