# ABclonal® www.abclonal.com

# **UXT Rabbit pAb**

Catalog No.: A4217

## **Basic Information**

#### **Observed MW**

17kDa

#### **Calculated MW**

18kDa

## Category

Polyclonal Antibody

## **Applications**

WB, ELISA

## **Cross-Reactivity**

Human

# **Background**

The protein encoded by this gene functions as a cofactor that modulates androgen receptor-dependent transcription, and also plays a critical role in tumor necrosis factor-induced apoptosis. Expression of this gene may play a role in tumorigenesis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

# **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**8409

Swiss Prot
Q9UBK9

## **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

# **Synonyms**

SKP2; STAP1; ART-27; UXT

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using UXT Rabbit pAb (A4217) at 1:1000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.