# ABclonal® www.abclonal.com

## 14-3-3 sigma Rabbit mAb

Catalog No.: A4377 Recombinant

## **Basic Information**

#### **Observed MW**

28kDa

#### **Calculated MW**

28kDa

#### Category

SMab Recombinant Monoclonal Antibody

## **Applications**

WB,IHC-P,ELISA

## **Cross-Reactivity**

Human, Mouse

#### CloneNo number

ARC0988

## **Background**

This gene encodes a cell cycle checkpoint protein. The encoded protein binds to translation and initiation factors and functions as a regulator of mitotic translation. In response to DNA damage this protein plays a role in preventing DNA errors during mitosis.

## **Recommended Dilutions**

WB 1:500 - 1:1000

**IHC-P** 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**2810

Swiss Prot
P31947

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

YWHAS; 14-3-3 sigma

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## **Validation Data**



Western blot analysis of various lysates using 14-3-3 sigma Rabbit mAb (A4377) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3min.



Immunohistochemistry analysis of paraffin-embedded Human colon using 14-3-3 sigma Rabbit mAb (A4377) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.