# ABclonal www.abclonal.com

# Cytokeratin 15 (KRT15) Rabbit mAb

Catalog No.: A4854 Recombinant 3 Publications

## **Basic Information**

## **Observed MW**

49kDa

#### **Calculated MW**

49kDa

#### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Rat

#### CloneNo number

ARC0270

# Background

The protein encoded by this gene is a member of the keratin gene family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. Most of the type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains and are clustered in a region on chromosome 17q21.2.

# **Recommended Dilutions**

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

Recommended starting **ELISA** 

> concentration is 1 µg/mL. Please optimize the concentration based on your specific

> > assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 3866 P19012

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

K15; CK15; K1CO; Cytokeratin 15 (KRT15)

# Contact

www.abclonal.com €

### **Product Information**


Source **Isotype Purification** Rabbit Affinity purification IgG

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# **Validation Data**



Western blot analysis of lysates from CAL 27 cells, using Cytokeratin 15 (KRT15) Rabbit mAb (A4854) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Immunofluorescence analysis of paraffin-embedded rat skin using Cytokeratin 15 (KRT15) (KRT15) Rabbit mAb (A4854) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.