# ABclonal®

## **SULT2A1 Rabbit pAb**

Catalog No.: A5489

## **Basic Information**

#### **Observed MW**

34kDa

#### **Calculated MW**

34kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse

## **Background**

This gene encodes a member of the sulfotransferase family. Sulfotransferases aid in the metabolism of drugs and endogenous compounds by converting these substances into more hydrophilic water-soluble sulfate conjugates that can be easily excreted. This protein catalyzes the sulfation of steroids and bile acids in the liver and adrenal glands, and may have a role in the inherited adrenal androgen excess in women with polycystic ovary syndrome.

## **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**6822

Swiss Prot
Q06520

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

HST; ST2; STD; hSTa; DHEAS; ST2A1; ST2A3; DHEA-ST; SULT2A3; DHEA-ST8; SULT2A1

## Contact

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using SULT2A1 Rabbit pAb (A5489) at 1:2000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.