

A5542

Leader in Biomolecular Solutions for Life Science



CGRP Rabbit pAb

Catalog No.: A5542

14 Publications

Basic Information

Observed MW

14kDa

Calculated MW

13kDa/15kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

This gene encodes the peptide hormones calcitonin, calcitonin gene-related peptide and katacalcin by tissue-specific alternative RNA splicing of the gene transcripts and cleavage of inactive precursor proteins. Calcitonin is involved in calcium regulation and acts to regulate phosphorus metabolism. Calcitonin gene-related peptide functions as a vasodilator and as an antimicrobial peptide while katacalcin is a calcium-lowering peptide. Multiple transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:1000 - 1:5000

IHC-P 1:2000 - 1:7000

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

796

Swiss Prot

P01258/P06881

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

CT; KC; PCT; CGRP; CALC1; CGRP1; CGRP-I; CGRP-alpha

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

Contact




www.abclonal.com


Validation Data




Western blot analysis of lysates from wild type (WT) and 293T cells transfected with CGRP using CGRP Rabbit pAb (A5542) at 1:2000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.




Immunohistochemistry analysis of paraffin-embedded Human thyroid medullary carcinoma tissue using CGRP Rabbit pAb (A5542) at a dilution of 1:6000 (40x lens). High pressure antigen retrieval performed with 0.01M Tris-EDTA Buffer(pH 9.0) prior to IHC staining.



Immunofluorescence analysis of C6 cells using CGRP Rabbit pAb (A5542) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using CGRP Rabbit pAb (A5542) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using CGRP Rabbit pAb (A5542) at dilution of 1:100. Blue: DAPI for nuclear staining.