# ABclonal www.abclonal.com

# FGG Rabbit pAb

Catalog No.: A5642 1 Publications

### **Basic Information**

### **Observed MW**

52kDa

#### **Calculated MW**

52kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse

## Background

The protein encoded by this gene is the gamma component of fibrinogen, a bloodborne glycoprotein comprised of three pairs of nonidentical polypeptide chains. Following vascular injury, fibrinogen is cleaved by thrombin to form fibrin which is the most abundant component of blood clots. In addition, various cleavage products of fibrinogen and fibrin regulate cell adhesion and spreading, display vasoconstrictor and chemotactic activities, and are mitogens for several cell types. Mutations in this gene lead to several disorders, including dysfibrinogenemia, hypofibrinogenemia and thrombophilia. Alternative splicing results in transcript variants encoding different isoforms.

# **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 2266 P02679

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

FGG

## **Contact**

www.abclonal.com €


### **Product Information**

Source **Isotype Purification** Rabbit Affinity purification IgG

### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of lysates from Mouse liver, using FGG Rabbit pAb (A5642) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).