# ABclonal® www.abclonal.com

## NOGO Receptor 1 (RTN4R) Rabbit pAb

Catalog No.: A5847 1 Publications

## **Basic Information**

#### **Observed MW**

66kDa

#### **Calculated MW**

51kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes the receptor for reticulon 4, oligodendrocyte myelin glycoprotein and myelin-associated glycoprotein. This receptor mediates axonal growth inhibition and may play a role in regulating axonal regeneration and plasticity in the adult central nervous system.

## **Recommended Dilutions**

WB 1:100 - 1:500

**IF/ICC** 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**65078
Swiss Prot
Q9BZR6

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

NGR; NOGOR; NOGO Receptor 1 (RTN4R)

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## **Validation Data**



Western blot analysis of various lysates using NOGO Receptor 1 (RTN4R) Rabbit pAb (A5847) at 1:500 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.



Immunofluorescence analysis of Neuro-2a cells using NOGO Receptor 1 (RTN4R) Rabbit pAb (A5847) at dilution of 1:20 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.