# ABclonal www.abclonal.com

## [KO Validated] SETDB1 Rabbit pAb

Catalog No.: A6145 KO Validated 2 Publications

### **Basic Information**

#### **Observed MW**

180kDa

#### **Calculated MW**

143kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

## Background

This gene encodes a histone methyltransferase which regulates histone methylation, gene silencing, and transcriptional repression. This gene has been identified as a target for treatment in Huntington Disease, given that gene silencing and transcription dysfunction likely play a role in the disease pathogenesis. Alternatively spliced transcript variants of this gene have been described.

## **Recommended Dilutions**

WB 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**9869

Swiss Prot
Q15047

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

ESET; KG1T; KMT1E; TDRD21; H3-K9-HMTase4; B1

### Contact

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from wild type (WT) and SETDB1 knockout (KO) 293T cells, using [KO Validated] SETDB1 Rabbit pAb (A6145) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Western blot analysis of various lysates using [KO Validated] SETDB1 Rabbit pAb (A6145) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.