# **SERPINA7 Rabbit pAb**

Catalog No.: A6313



## **Basic Information**

#### **Observed MW**

46kDa

### **Calculated MW**

46kDa

### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

### **Background**

There are three proteins including thyroxine-binding globulin (TBG), transthyretin and albumin responsible for carrying the thyroid hormones thyroxine (T4) and 3,5,3'-triiodothyronine (T3) in the bloodstream. This gene encodes the major thyroid hormone transport protein, TBG, in serum. It belongs to the serpin family in genomics, but the protein has no inhibitory function like many other members of the serpin family. Mutations in this gene result in TGB deficiency, which has been classified as partial deficiency, complete deficiency, and excess, based on the level of serum TBG. Alternatively spliced transcript variants encoding different isoforms have been found, but the full-length nature of these variants has not been determined.

# **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

### **Immunogen Information**

**Gene ID**6906

Swiss Prot
P05543

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

TBG; TBGQTL; SERPINA7

### **Contact**

www.abclonal.com


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using SERPINA7 Rabbit pAb (A6313) at 1:1000 dilution.\_Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.\_Lysates/proteins: 25 $\mu$ g per lane.\_Blocking buffer: 3% nonfat dry milk in TBST.\_Detection: ECL Enhanced Kit (RM00021).\_Exposure time: 30s.