# SPINT1 Rabbit pAb

Catalog No.: A6411



### **Basic Information**

#### **Observed MW**

70kDa

#### **Calculated MW**

58kDa

#### **Category**

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene is a member of the Kunitz family of serine protease inhibitors. The protein is a potent inhibitor specific for HGF activator and is thought to be involved in the regulation of the proteolytic activation of HGF in injured tissues. Alternative splicing results in multiple variants encoding different isoforms.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:100

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

# **Immunogen Information**

**Gene ID**Swiss Prot
6692
043278

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

# **Synonyms**

HAI; HAI1; MANSC2; SPINT1

## Contact

www.abclonal.com


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification


#### Storage

Store at -20 $^{\circ}$ C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of various lysates using SPINT1 Rabbit pAb (A6411) at 1:1000 dilution.\_Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.\_Lysates/proteins:  $25\mu g$  per lane.\_Blocking buffer: 3% nonfat dry milk in TBST.\_Detection: ECL Enhanced Kit (RM00021).\_Exposure time: 90s.



Immunofluorescence analysis of HeLa cells using SPINT1 Rabbit pAb (A6411). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.