# ABclonal www.abclonal.com

# MARK2 Rabbit mAb

Catalog No.: A6512 Recombinant

## **Basic Information**

## **Observed MW**

78kDa/82kDa

#### **Calculated MW**

88kDa

#### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC1862

# Background

This gene encodes a member of the Par-1 family of serine/threonine protein kinases. The protein is an important regulator of cell polarity in epithelial and neuronal cells, and also controls the stability of microtubules through phosphorylation and inactivation of several microtubule-associating proteins. The protein localizes to cell membranes. Multiple transcript variants encoding different isoforms have been found for this gene.

# **Recommended Dilutions**

WB 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 2011 Q7KZI7

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

EMK1; EMK-1; PAR-1; Par1b; Par-1b; MARK2

## Contact

www.abclonal.com €

### **Product Information**


Source **Isotype Purification** Rabbit Affinity purification IgG

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

# Validation Data



Western blot analysis of various lysates using MARK2 Rabbit mAb (A6512) at 1:1000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.