# ABclonal® www.abclonal.com

## **GLP2R Rabbit pAb**

Catalog No.: A6602 1 Publications

## **Basic Information**

#### **Observed MW**

63kDa

#### **Calculated MW**

63kDa

#### **Category**

Polyclonal Antibody

## **Applications**

WB,IHC-P,ELISA

## **Cross-Reactivity**

Human, Mouse

## **Background**

This gene encodes a G protein-coupled receptor that is closely related to the glucagon receptor and binds to glucagon-like peptide-2 (GLP2). Signalling through GLP2 stimulates intestinal growth and increases villus height in the small intestine, concomitant with increased crypt cell proliferation and decreased enterocyte apoptosis.

## **Recommended Dilutions**

WB 1:100 - 1:500

**IHC-P** 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**9340

Swiss Prot
9340

O95838

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

GLP2R

## **Contact**


www.abclonal.com

## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of various lysates using GLP2R Rabbit pAb (A6602) at 1:500 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Human colon using GLP2R Rabbit pAb (A6602) at dilution of 1:20 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.