# ABclonal www.abclonal.com

# CDKN1C/p57 Kip2 Rabbit mAb

Catalog No.: A6843 Recombinant 1 Publications

### **Basic Information**

#### **Observed MW**

57kDa

#### **Calculated MW**

32kDa

### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human

#### CloneNo number

ARC1424

# Background

This gene is imprinted, with preferential expression of the maternal allele. The encoded protein is a tight-binding, strong inhibitor of several G1 cyclin/Cdk complexes and a negative regulator of cell proliferation. Mutations in this gene are implicated in sporadic cancers and Beckwith-Wiedemann syndorome, suggesting that this gene is a tumor suppressor candidate. Three transcript variants encoding two different isoforms have been found for this gene.

# **Recommended Dilutions**

WB 1:2000 - 1:8000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID Swiss Prot** 1028 P49918

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

BWS; WBS; p57; BWCR; KIP2; p57Kip2; CDKN1C/p57 Kip2

### **Contact**

€ www.abclonal.com


## **Product Information**

Source **Isotype Purification** Rabbit Affinity purification IgG

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## **Validation Data**



Western blot analysis of lysates from HeLa cells using CDKN1C/p57 Kip2 Rabbit mAb (A6843) at 1:2000 dilution incubated overnight at 4°C. HeLa cells were treated with Dexamethasone (500 nM) at 37°C for 16hrs .

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3 % nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.