# ABclonal® www.abclonal.com

## MAPK13 Rabbit pAb

Catalog No.: A7496

## **Basic Information**

#### **Observed MW**

42kDa

#### **Calculated MW**

42kDa

### Category

Polyclonal Antibody

#### **Applications**

WB,ELISA

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a member of the mitogen-activated protein (MAP) kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. The encoded protein is a p38 MAP kinase and is activated by proinflammatory cytokines and cellular stress. Substrates of the encoded protein include the transcription factor ATF2 and the microtubule dynamics regulator stathmin. Alternatively spliced transcript variants have been observed for this gene.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
5603
O15264

## **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

SAPK4; PRKM13; MAPK 13; MAPK-13; p38delta; MAPK13

## **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using MAPK13 Rabbit pAb (A7496) at 1:1000 dilution.\_Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.\_Lysates/proteins: 25 $\mu$ g per lane.\_Blocking buffer: 3% nonfat dry milk in TBST.\_Detection: ECL Enhanced Kit (RM00021).\_Exposure time: 90s.