# ABclonal® www.abclonal.com

## ATF7 Rabbit pAb

Catalog No.: A7580

## **Basic Information**

#### **Observed MW**

53kDa

#### **Calculated MW**

52kDa

#### Category

Polyclonal Antibody

## **Applications**

WB, ELISA

## **Cross-Reactivity**

Human

## **Background**

Enables several functions, including DNA-binding transcription repressor activity, RNA polymerase II-specific; mitogen-activated protein kinase binding activity; and transcription coactivator binding activity. Involved in negative regulation of transcription by RNA polymerase II and positive regulation of transcription by RNA polymerase II. Located in nucleoplasm. Biomarker of colorectal cancer.

## **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
11016
P17544

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

#### **Synonyms**

ATFA; ATF7

## Contact

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using ATF7 Rabbit pAb (A7580) at 1:1000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.