

A7737

Leader in Biomolecular Solutions for Life Science



SYT4 Rabbit pAb

Catalog No.: A7737

1 Publications

Basic Information

Observed MW

46kDa

Calculated MW

48kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Human,Mouse,Rat

Background

Predicted to enable several functions, including calcium ion binding activity; phospholipid binding activity; and syntaxin binding activity. Involved in negative regulation of catecholamine secretion and positive regulation of dendrite extension. Predicted to be located in several cellular components, including microvesicle; perinuclear region of cytoplasm; and secretory vesicle. Predicted to be active in several cellular components, including axon; exocytic vesicle; and glutamatergic synapse. Predicted to be integral component of neuronal dense core vesicle membrane.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

IF/ICC 1:50 - 1:200

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

6860

Swiss Prot

Q9H2B2

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

HsT1192; SYT4

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Contact




www.abclonal.com


Validation Data




Western blot analysis of various lysates, using SYT4 Rabbit pAb (A7737) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Human stomach using SYT4 Rabbit pAb (A7737) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human liver cancer using SYT4 Rabbit pAb (A7737) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of U2OS cells using SYT4 Rabbit pAb (A7737) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.