# ABclonal www.abclonal.com

## **HSPA2** Rabbit pAb

Catalog No.: A7902 1 Publications

## **Basic Information**

#### **Observed MW**

75kDa

#### **Calculated MW**

70kDa

#### Category

Polyclonal Antibody

## **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Enables disordered domain specific binding activity; enzyme binding activity; and unfolded protein binding activity. Involved in negative regulation of inclusion body assembly and protein refolding. Located in cytosol.

## **Recommended Dilutions**

**WB** 1:500 - 1:2000

**IF/ICC** 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## 3306

**Gene ID** 

**Immunogen**Recombinant protein (or fragment). This information is considered to be commercially sensitive.

**Swiss Prot** 

P54652

#### **Synonyms**

HSP70-2; HSP70-3; HSPA2

**Immunogen Information** 

## **Contact**

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using HSPA2 Rabbit pAb (A7902) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.



Immunofluorescence analysis of L929 cells using HSPA2 Rabbit pAb (A7902) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using HSPA2 Rabbit pAb (A7902) at dilution of 1:100. Blue: DAPI for nuclear staining.