# **AURKC Rabbit pAb**

Catalog No.: A7930 1 Publications



# **Basic Information**

#### **Observed MW**

36kDa

#### **Calculated MW**

36kDa

#### **Category**

Polyclonal Antibody

### **Applications**

WB,IHC-P,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a member of the Aurora subfamily of serine/threonine protein kinases. The encoded protein is a chromosomal passenger protein that forms complexes with Aurora-B and inner centromere proteins and may play a role in organizing microtubules in relation to centrosome/spindle function during mitosis. This gene is overexpressed in several cancer cell lines, suggesting an involvement in oncogenic signal transduction. Alternative splicing results in multiple transcript variants.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

IHC-P 1:50 - 1:100

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

# **Immunogen Information**

**Gene ID** Swiss Prot 6795 Q9UQB9

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

AIE2; AIK3; ARK3; AurC; SPGF5; STK13; HEL-S-90; aurora-C; AURKC

### Contact

www.abclonal.com

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using AURKC Rabbit pAb (A7930) at 1:500

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.