ABclonal www.abclonal.com

RRM2B Rabbit pAb

Catalog No.: A8020 1 Publications

Basic Information

Observed MW

Refer to figures

Calculated MW

41kDa

Category

Polyclonal Antibody

Applications

IF/ICC,ELISA

Cross-Reactivity

Human

Background

This gene encodes the small subunit of a p53-inducible ribonucleotide reductase. This heterotetrameric enzyme catalyzes the conversion of ribonucleoside diphosphates to deoxyribonucleoside diphosphates. The product of this reaction is necessary for DNA synthesis. Mutations in this gene have been associated with autosomal recessive mitochondrial DNA depletion syndrome, autosomal dominant progressive external ophthalmoplegia-5, and mitochondrial neurogastrointestinal encephalopathy. Alternatively spliced transcript variants have been described.

Recommended Dilutions

IF/ICC 1:50 - 1:200

ELISA

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID50484

Swiss Prot
Q7LG56

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

P53R2; RCDFRD; MTDPS8A; MTDPS8B; RRM2B

Contact

www.abclonal.com


Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data





Immunofluorescence analysis of U-2 OS cells using RRM2B Rabbit pAb (A8020) at a dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of HeLa cells using RRM2B Rabbit pAb (A8020) at a dilution of 1:200 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.