# ABclonal www.abclonal.com

# **Gephyrin Rabbit pAb**

Catalog No.: A8572 2 Publications

### **Basic Information**

### **Observed MW**

103kDa

#### **Calculated MW**

80kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

## Background

This gene encodes a neuronal assembly protein that anchors inhibitory neurotransmitter receptors to the postsynaptic cytoskeleton via high affinity binding to a receptor subunit domain and tubulin dimers. In nonneuronal tissues, the encoded protein is also required for molybdenum cofactor biosynthesis. Mutations in this gene may be associated with the neurological condition hyperplexia and also lead to molybdenum cofactor deficiency. Numerous alternatively spliced transcript variants encoding different isoforms have been described; however, the full-length nature of all transcript variants is not currently known.

# **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID Swiss Prot** 10243 Q9NQX3

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

## **Synonyms**

GPH; GEPH; HKPX1; GPHRYN; MOCODC; Gephyrin

### **Contact**

www.abclonal.com €


### **Product Information**

Source **Isotype Purification** Rabbit Affinity purification IgG

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of various lysates using Gephyrin Rabbit pAb (A8572) at 1:1000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000  $\,$ 

dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.