ABclonal

www.abclonal.com

Sec23A Rabbit mAb

Catalog No.: A8613 Recombinant

Basic Information

Observed MW

85kDa

Calculated MW

86kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IF/ICC,ELISA

Cross-Reactivity

Human, Mouse

CloneNo number

ARC1754

Background

The protein encoded by this gene is a member of the SEC23 subfamily of the SEC23/SEC24 family. It is part of a protein complex and found in the ribosome-free transitional face of the endoplasmic reticulum (ER) and associated vesicles. This protein has similarity to yeast Sec23p component of COPII. COPII is the coat protein complex responsible for vesicle budding from the ER. The encoded protein is suggested to play a role in the ER-Golgi protein trafficking.

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

Recommended starting **ELISA**

> concentration is 1 µg/mL. Please optimize the concentration based on your specific

> > assay requirements.

Immunogen Information

Gene ID Swiss Prot 10484 Q15436

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CLSD; hSec23A; Sec23A

Contact

www.abclonal.com €

Product Information


Source **Isotype Purification** Rabbit Affinity purification IgG

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from Hep G2 cells, using Sec23AA Rabbit mAb (A8613) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.



Immunofluorescence analysis of NIH/3T3 cells using Sec23AA Rabbit mAb (A8613) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.