# CD74 Rabbit mAb

Catalog No.: A9149 Recombinant



## **Basic Information**

### **Observed MW**

30-40kDa

### **Calculated MW**

34kDa

### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB,IHC-P,ELISA

### **Cross-Reactivity**

Human

### CloneNo number

ARC1452

## **Background**

The protein encoded by this gene associates with class II major histocompatibility complex (MHC) and is an important chaperone that regulates antigen presentation for immune response. It also serves as cell surface receptor for the cytokine macrophage migration inhibitory factor (MIF) which, when bound to the encoded protein, initiates survival pathways and cell proliferation. This protein also interacts with amyloid precursor protein (APP) and suppresses the production of amyloid beta (Abeta). Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

## **Recommended Dilutions**

**WB** 1:500 - 1:1000

**IHC-P** 1:50 - 1:200

**ELISA** Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

## **Immunogen Information**

**Gene ID**972

Swiss Prot
P04233

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

II; p33; CLIP; DHLAG; HLADG; Ia-GAMMA; CD74

## Contact

www.abclonal.com


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.



Western blot analysis of various lysates using CD74 Rabbit mAb (A9149) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using CD74 Rabbit mAb (A9149) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human small cell lung cancer using CD74 Rabbit mAb (A9149) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human tonsil using CD74 Rabbit mAb (A9149) at dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.