

A9377

Leader in Biomolecular Solutions for Life Science



SR protein repeat Rabbit mAb

Catalog No.: A9377

Recombinant

Basic Information

Observed MW

40kDa

Calculated MW

40kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB,IHC-P,IP,ELISA

Cross-Reactivity

Human,Mouse,Rat

CloneNo number

ARC2754

Background

The protein encoded by this gene is involved in mRNA splicing and may play a role in the determination of alternative splicing. The encoded nuclear protein belongs to the splicing factor SR family and has been shown to bind with and modulate another member of the family, SFRS12. Alternative splicing results in multiple transcript variants. In addition, two pseudogenes, one on chromosome 17 and the other on the X chromosome, have been found for this gene.

Recommended Dilutions

WB 1:100 - 1:500

IHC-P 1:50 - 1:200

IP 0.5µg-4µg antibody for
300µg-600µg extracts
of whole cells

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Immunogen Information

Gene ID

6431

Swiss Prot

Q13247

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

B52; SFRS6; SRP55; HEL-S-91; SR protein repeat

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

Contact




www.abclonal.com


Validation Data




Western blot analysis of various lysates using (A9377) at 1:500 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 30s.




Western blot analysis of lysates from Rat liver, using (A9377) at 1:500 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Enhanced Kit (RM00021).
 Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Human colon tissue using SR protein repeat Rabbit mAb (A9377) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Rat colon tissue using SR protein repeat Rabbit mAb (A9377) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human breast tissue using SR protein repeat Rabbit mAb (A9377) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma tissue using SR protein repeat Rabbit mAb (A9377) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M Tris-EDTA buffer (pH 9.0) prior to IHC staining.



Immunoprecipitation of SR protein repeat in 600 μ g extracts from Mouse thymus using 3 μ g SR protein repeat Rabbit mAb (A9377). Western blot analysis was performed using SR protein repeat Rabbit mAb (A9377) at 1:500 dilution.