

A9712

Leader in Biomolecular Solutions for Life Science



EAAT1/SLC1A3 Rabbit mAb

Catalog No.: A9712

Recombinant

Basic Information

Observed MW

50-55kDa/90-130kDa

Calculated MW

60kDa

Category

SMab Recombinant Monoclonal
Antibody

Applications

WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity

Mouse,Rat

CloneNo number

ARC1714

Recommended Dilutions

WB 1:500 - 1:2000

IHC-P 1:200 - 1:2000

IF/ICC 1:200 - 1:800

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Contact



www.abclonal.com

Background

This gene encodes a member of a member of a high affinity glutamate transporter family. This gene functions in the termination of excitatory neurotransmission in central nervous system. Mutations are associated with episodic ataxia, Type 6. Alternative splicing results in multiple transcript variants.

Immunogen Information

Gene ID

6507

Swiss Prot

P43003

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

EA6; EAAT1; GLAST; GLAST1; EAAT1/SLC1A3

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


Validation Data




Western blot analysis of various lysates using EAAT1/SLC1A3 Rabbit mAb (A9712) at 1:1000 dilution incubated overnight at 4°C.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 45s.




Immunohistochemistry analysis of paraffin-embedded Mouse brain tissue using EAAT1/SLC1A3 Rabbit mAb (A9712) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat brain tissue using EAAT1/SLC1A3 Rabbit mAb (A9712) at a dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Confocal imaging of paraffin-embedded Rat brain tissue using EAAT1/SLC1A3 Rabbit mAb (A9712, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform microwave antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.



Confocal imaging of paraffin-embedded Mouse brain tissue using EAAT1/SLC1A3 Rabbit mAb (A9712, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used

Validation Data

for nuclear staining (Blue). Objective:
40x. Perform microwave antigen
retrieval with 0.01 M citrate buffer
(pH 6.0) prior to IF staining.