

A9713

Leader in Biomolecular Solutions for Life Science



UACA Rabbit pAb

Catalog No.: A9713

Basic Information

Observed MW

162kDa

Calculated MW

163kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a protein that contains ankyrin repeats and coiled coil domains and likely plays a role in apoptosis. Studies in rodents have implicated the encoded protein in the stimulation of apoptosis and the regulation of mammary gland involution, in which the mammary gland returns to its pre-pregnant state. This protein has also been proposed to negatively regulate apoptosis based on experiments in human cell lines in which the protein was shown to interact with PRKC apoptosis WT1 regulator protein, also known as PAR-4, and inhibit translocation of the PAR-4 receptor. Autoantibodies to this protein have been identified in human patients with panuveitis and Graves' disease. Differential expression of this gene has been observed in various human cancers.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID

55075

Swiss Prot

Q9BZF9

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

NUCLING; UACA

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

Validation Data



Western blot analysis of various lysates using UACA Rabbit pAb (A9713) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 5s.