# JAM2 Rabbit pAb

Catalog No.: A9978 2 Publications



### **Basic Information**

### **Observed MW**

36kDa

### **Calculated MW**

33kDa

### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# Background

This gene belongs to the immunoglobulin superfamily, and the junctional adhesion molecule (JAM) family. The protein encoded by this gene is a type I membrane protein that is localized at the tight junctions of both epithelial and endothelial cells. It acts as an adhesive ligand for interacting with a variety of immune cell types, and may play a role in lymphocyte homing to secondary lymphoid organs. Alternatively spliced transcript variants have been found for this gene.

# **Recommended Dilutions**

WB 1:500 - 1:1000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

# **Immunogen Information**

**Gene ID**58494

Swiss Prot
P57087

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

JAMB; CD322; IBGC8; JAM-B; VEJAM; PRO245; VE-JAM; C21orf43; JAM2

### **Contact**

www.abclonal.com


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using JAM2 Rabbit pAb (A9978) at 1:1000

dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000


dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Western blot analysis of various lysates using JAM2 Rabbit pAb (A9978) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.