

AE095

Leader in Biomolecular Solutions for Life Science



# HRP-conjugated Rabbit anti DDDDK-Tag mAb

Catalog No.: AE095

1 Publications

## Basic Information

### Observed MW

50kDa/46kDa

### Calculated MW

### Category

SMab Recombinant Monoclonal  
Antibody

### Applications

WB, ELISA

### Cross-Reactivity

Species independent

### CloneNo number

ARC5111-01-HRP

### Conjugate

HRP

## Recommended Dilutions

**WB** 1:2000-6000

**ELISA** Recommended starting  
concentration is 1  
µg/mL. Please optimize  
the concentration  
based on your specific  
assay requirements.

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Background

FLAG-tag, or FLAG octapeptide, or FLAG epitope, is a polypeptide protein tag that can be added to a protein using recombinant DNA technology, having the sequence motif DYKDDDDK. It has been used for studying proteins in living cells and for protein purification by affinity chromatography. It has been used to separate recombinant, overexpressed protein from wild-type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits, because its mild purification procedure tends not to disrupt such complexes. It has been used to obtain proteins of sufficient purity and quality to carry out 3D structure determination by x-ray crystallography. A FLAG-tag can be used in many different assays that require recognition by an antibody. If there is no antibody against a given protein, adding a FLAG-tag to a protein allows the protein to be studied with an antibody against the FLAG sequence. Examples are cellular localization studies by immunofluorescence or detection by SDS PAGE protein electrophoresis and Western blotting.

## Immunogen Information

### Gene ID

### Swiss Prot

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

DDDDK; DDDDK tag; DDDDK-tag

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Western blot analysis of extracts of normal 293T cells, 293T transfected with GSK3B-C Protein, using HRP-conjugated rabbit anti DDDDK-Tag mAb (AE095) at 1:5000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Western blot analysis of extracts of normal 293F , 293F transfected with IRF1-N Protein, using HRP-conjugated rabbit anti DDDDK-Tag mAb (AE095) at 1:5000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.