# mCherry Rabbit pAb

Catalog No.: AE171 2 Publications



#### **Basic Information**

**Observed MW** 

55kDa

**Calculated MW** 

**Category** 

Polyclonal Antibody

**Applications** 

WB,ELISA

**Cross-Reactivity** 

Species independent

## **Background**

Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.

#### **Recommended Dilutions**

WB 1:2000 - 1:10000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

Gene ID Swiss Prot

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

**Synonyms** 

mCherry; mCherry tag; mCherry-tag

#### Contact

€

www.abclonal.com


#### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from wild type (WT) and 293F cells transfected with CT55-mCherry using mCherry Rabbit pAb (AE171) at 1:10000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.