# ABclonal www.abclonal.com

## Phospho-RB-S811 Rabbit pAb

Catalog No.: AP0089 3 Publications

## **Basic Information**

#### **Observed MW**

106kDa

#### **Calculated MW**

106kDa

#### **Category**

Polyclonal Antibody

## **Applications**

WB, ELISA

## **Cross-Reactivity**

Human

## **Background**

The protein encoded by this gene is a negative regulator of the cell cycle and was the first tumor suppressor gene found. The encoded protein also stabilizes constitutive heterochromatin to maintain the overall chromatin structure. The active, hypophosphorylated form of the protein binds transcription factor E2F1. Defects in this gene are a cause of childhood cancer retinoblastoma (RB), bladder cancer, and osteogenic sarcoma.

## **Recommended Dilutions**

**WB** 1:1000 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**5925

Swiss Prot
P06400

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

### **Synonyms**

RB; pRb; OSRC; pp110; p105-Rb; PPP1R130; p110-RB1; Phospho-RB-S811

## Contact

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from HeLa cells, using Phospho-RB-S811 Rabbit pAb (AP0089) at 1:1000 dilution. HeLa cells were treated with UV for 15-30 minutes. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% BSA.