# ABclonal®

## Phospho-Stathmin 1-S16 Rabbit pAb

Catalog No.: AP0229

## **Basic Information**

#### **Observed MW**

15kDa

#### **Calculated MW**

17kDa

## **Category**

Polyclonal Antibody

#### **Applications**

WB, ELISA

## **Cross-Reactivity**

Human, Mouse

## **Background**

This gene belongs to the stathmin family of genes. It encodes a ubiquitous cytosolic phosphoprotein proposed to function as an intracellular relay integrating regulatory signals of the cellular environment. The encoded protein is involved in the regulation of the microtubule filament system by destabilizing microtubules. It prevents assembly and promotes disassembly of microtubules. Multiple transcript variants encoding different isoforms have been found for this gene.

## **Recommended Dilutions**

WB 1:500 - 1:2000

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**Swiss Prot
3925
P16949

#### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

#### **Synonyms**

Lag; SMN; OP18; PP17; PP19; PR22; LAP18; C1orf215; Phospho-Stathmin 1-S16

## Contact

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from HeLa and C2C12 cells, using Phospho-Stathmin 1-S16 Rabbit pAb (AP0229) at 1:2000 dilution. HeLa and C2C12 cells were treated with Nocodazole (50ng/mL) for 20 hours.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.