

AP0417

Leader in Biomolecular Solutions for Life Science



# Phospho-NFKB1-S932 Rabbit pAb

Catalog No.: AP0417

2 Publications

## Basic Information

**Observed MW**

105kDa

**Calculated MW**

105kDa

**Category**

Polyclonal Antibody

**Applications**

WB

**Cross-Reactivity**

Human,Mouse,Rat

## Background

This gene encodes a 105 kD protein which can undergo cotranslational processing by the 26S proteasome to produce a 50 kD protein. The 105 kD protein is a Rel protein-specific transcription inhibitor and the 50 kD protein is a DNA binding subunit of the NF-kappa-B (NFKB) protein complex. NFKB is a transcription regulator that is activated by various intra- and extra-cellular stimuli such as cytokines, oxidant-free radicals, ultraviolet irradiation, and bacterial or viral products. Activated NFKB translocates into the nucleus and stimulates the expression of genes involved in a wide variety of biological functions. Inappropriate activation of NFKB has been associated with a number of inflammatory diseases while persistent inhibition of NFKB leads to inappropriate immune cell development or delayed cell growth. NFKB is a critical regulator of the immediate-early response to viral infection. Alternative splicing results in multiple transcript variants encoding different isoforms, at least one of which is proteolytically processed.

## Recommended Dilutions

WB

1:500 - 1:2000

## Immunogen Information

**Gene ID**

4790

**Swiss Prot**

P19838

**Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

**Synonyms**

KBF1; EBP-1; NF-kB; CVID12; NF-kB1; NFKB-p50; NFkappaB; NF-kappaB; NFKB-p105; NF-kappa-B1; NF-kappabeta; Phospho-NFKB1-S932

## Contact

[www.abclonal.com](http://www.abclonal.com)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of lysates from 293 cells, using Phospho-NFKB1-S932 Rabbit pAb (AP0417). 293 cells treated with TNF-α.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25μg per lane.

Blocking buffer: 3% BSA.