

AP0467

Leader in Biomolecular Solutions for Life Science



Phospho-ZAP70-Y319 Rabbit pAb

Catalog No.: AP0467

Basic Information

Observed MW

70kDa

Calculated MW

70kDa

Category

Polyclonal Antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Human

Background

This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:2000

IP 0.5µg-4µg antibody for
200µg-400µg extracts
of whole cells

ELISA Recommended starting
concentration is 1
µg/mL. Please optimize
the concentration
based on your specific
assay requirements.

Contact



www.abclonal.com

Immunogen Information

Gene ID

7535

Swiss Prot

P43403

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

SRK; STD; TZK; STCD; IMD48; ADMIO2; ZAP-70; Phospho-ZAP70-Y319

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Immunoprecipitation analysis of 200 µg extracts of Jurkat cells, using 3 µg Phospho-ZAP70-Y319 pAb (AP0467). Western blot was performed from the immunoprecipitate using Phospho-ZAP70-Y319 pAb (AP0467) at a dilution of 1:1000. Jurkat cells were treated with Hydrogen Peroxide (2 mM) at 37°C for 2 minutes after serum-starvation overnight.