

AP0492

Leader in Biomolecular Solutions for Life Science



Phospho-TrkA-Y490 Rabbit pAb

Catalog No.: AP0492

2 Publications

Basic Information

Observed MW

150kDa/140kDa

Calculated MW

87kDa

Category

Polyclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the neurotrophic tyrosine kinase receptor (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. The presence of this kinase leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in this gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, cognitive disability and cancer. Alternate transcriptional splice variants of this gene have been found, but only three have been characterized to date.

Recommended Dilutions

WB 1:500 - 1:2000

ELISA Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID
4914

Swiss Prot
P04629

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

MTC; TRK; TRK1; TRKA; Trk-A; p140-TrkA; Phospho-TrkA-Y490

Contact

 www.abclonal.com

Product Information

Source
Rabbit

Isotype
IgG


Purification
Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data




Western blot analysis of lysates from NIH3T3 cells, using Phospho-TrkA-Y490 Rabbit pAb (AP0492).

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.



Western blot analysis of lysates from PC-12 cells using Phospho-TrkA-Y490 Rabbit pAb (AP0492) at 1:1800 dilution. PC-12 cells were treated by MNGF (200 ng/ml) at 37°C for 10 minutes after serum-starvation overnight.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 30 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.