

AP0524

Leader in Biomolecular Solutions for Life Science



# Phospho- $\beta$ -Catenin-S33/S37/T41 Rabbit pAb

Catalog No.: AP0524

1 Publications

## Basic Information

### Observed MW

Refer to figures

### Calculated MW

85kDa

### Category

Polyclonal Antibody

### Applications

IHC-P, ELISA

### Cross-Reactivity

Human, Mouse, Rat

## Background

The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript variants.

## Recommended Dilutions

IHC-P 1:50 - 1:200

ELISA Recommended starting concentration is 1  $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

### Gene ID

1499

### Swiss Prot

P35222

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

EVR7; CTNNB; MRD19; NEDSDV; armadillo; Phospho- $\beta$ -Catenin-S33/S37/T41

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification


### Storage

Store at -20°C. Avoid freeze / thaw cycles.


Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Validation Data


---



Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using Phospho-β-Catenin-S33/S37/T41 Rabbit pAb (AP0524) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat skin using Phospho-β-Catenin-S33/S37/T41 Rabbit pAb (AP0524) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney using Phospho-β-Catenin-S33/S37/T41 Rabbit pAb (AP0524) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M Tris/EDTA Buffer (pH 9.0) prior to IHC staining.