# ABclonal www.abclonal.com

## Phospho-SIRT1-S47 Rabbit mAb

Catalog No.: AP0976 Recombinant 2 Publications

## **Basic Information**

## **Observed MW**

120kDa

#### **Calculated MW**

82kDa

#### Category

SMab Recombinant Monoclonal Antibody

## **Applications**

WB, ELISA

## **Cross-Reactivity**

Human

#### CloneNo number

ARC1520

## **Background**

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Alternative splicing results in multiple transcript variants.

## **Recommended Dilutions**

**WB** 1:100 - 1:500

**ELISA** 

Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## **Immunogen Information**

**Gene ID**23411

Swiss Prot
Q96EB6

### **Immunogen**

Synthetic peptide. This information is considered to be commercially sensitive.

## **Synonyms**

SIR2; SIR2L1; SIR2alpha; Phospho-SIRT1-S47

## Contact

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20℃. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of lysates from 293T cells, using Phospho-SIRT1-S47 Rabbit mAb (AP0976) at 1:500 dilution. 293T cells were treated by CIP(20uL/400ul) at 37°C for 1

hour.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 180s.