ABclonal www.abclonal.com

Pan Phospho-Tyrosine Rabbit mAb

Catalog No.: AP1162 Recombinant 3 Publications

Basic Information

Observed MW

Calculated MW

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IP,ELISA

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC54383

Background

Tyrosine phosphorylation (pTyr), much of which occurred on localized multiple sites, initiates cellular signaling, governs cellular functions, and its dysregulation is implicated in many diseases, especially cancers. pTyr-specific sensing is of great significance for understanding disease states and developing targeted anticancer drugs.

Recommended Dilutions

WB 1:500 - 1:1000

IP 0.5μg-4μg antibody for

200μg-400μg extracts

of whole cells

ELISA Recommended starting

concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Contact

www.abclonal.com

Immunogen Information

Gene ID Swiss Prot

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.09% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates using Pan Phospho-Tyrosine Rabbit mAb (AP1162) at 1:1000 dilution. Jurkat cells and NIH/3T3 cells and C6 cells were treated with Pervanadate (1 mM) at 37° C for 30 minutes.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% BSA.

Detection: ECL Basic Kit (RM00020).

Exposure time: 5s.



Immunoprecipitation analysis of 300 μ g extracts of Jurkat (pervanadate) cells using 3 μ g Pan Phospho-Tyrosine Rabbit mAb (AP1162). Western blot was performed from the immunoprecipitate using Pan Phospho-Tyrosine Rabbit mAb (AP1162) at a dilution of 1:1000.